Complete if Known Substitute for form 1449/PTO Application Number 10/586,015 INFORMATION DISCLOSURE Filing Date August 4, 2008 STATEMENT BY APPLICANT First Named Inventor Eric T. Ahrens Art Unit 1618 (Use as many sheets as necessary) Jagadishwar Rao Samala Examiner Name 4 CAMU-P01-002 1 of Sheet Attorney Docket Number

	U.S. PATENT DOCUMENTS				
Examiner	Cite	Document Number	Publication Date	Name of Patentee or	Pages, Columns, Lines, Where
Initials*	No.1	Number-Kind Code <sup>2</sup> ( if known)	MM-DD-YYYY	MM DD.VVV Applicant of Cited Decument	Relevant Passages or Relevant Figures Appear
	AT1	2002/0192688	12-19-2002	Yang et al.	
	AU1	2004/0109824	06-10-2004	Hinds et al.	
	AV1	2005/0008572	01-13-2005	Prokop et al.	
	AW1	2005/0244384	11-03-2005	Law	
	AX1	2006/0040389	02-23-2006	Murry et al	
	AY1	2006/0239919	10-26-2006	Wickline et al.	
	AX1	2007/0253910	11-01-2007	Ahrens et al.	
	AA2	2007/0258886	11-18-2007	Ahrens et al.	
	AB2	2009/0263329	10-22-2009	Wickline et al.	
	AC2	4,094,911	06-13-1978	Zollinger	
	AD2	5,785,950	07-28-1998	Kaufman et al.	
	AE2	5,958,371	09-28-1999	Lanza et al.	
	AF2	7,357,937	04-15-2008	Hsu et al.	
	AG2	7,514,074	04-07-2009	Pittinger et al.	

	FOREIGN PATENT DOCUMENTS						
Examiner Initials*	Cite No.1	Foreign Patent Document  Country Code <sup>3</sup> -Number <sup>4</sup> -Kind Code <sup>5</sup> (if known)	Publication Date MM-DD-YYYY	Name of Patentee or Applicant of Cited Document	Pages, Columns, Lines, Where Relevant Passages Or Relevant Figures Appear		
	BK	EP 1 728 788	12-06-2006	Shinetsu Chemical Co.			
	BL	WO91/14664	10-03-1991	Hider et al.			
	ВМ	WO94/18954	09-01-1994	Clover Consolidated, Limited			
	BN	WO96/41647	12-27-1996	Barnes-Jewish Hospital			
	ВО	WO97/40679	11-06-1997	Imarz Pharmaceuticals Corp.			
	BP	WO05/072780	08-11-2005	Carnegie Mellon University			
	BQ	WO06/096499	09-14-2006	Washington University			
	BR	WO07/100715	09-07-2007	Washington University			
	BS	WO08/119790	10-09-2008	Heinrich-Heine Universität Düsseldorf			
	BT	WO09/009105	01-15-2009	Carnegie Mellon University			

\*EXAMINER: Initial if reference considered, whether or not citation is in conformance with MPEP 609. Draw line through citation if not in conformance and not considered. Include copy of this form with next communication to applicant. <sup>1</sup> Applicant's unique citation designation number (optional). <sup>2</sup> See Kinds Codes of USPTO Patent Documents at <a href="https://www.uspto.gov">www.uspto.gov</a> or MPEP 901.04. <sup>3</sup> Enter Office that issued the document, by the two-letter code (WIPO Standard ST.3). <sup>4</sup> For Japanese patent documents, the indication of the year of the reign of the Emperor must precede the serial number of the patent document. <sup>5</sup> Kind of document by the appropriate symbols as indicated on the document under WIPO Standard ST.16 if possible. <sup>6</sup> Applicant is to place a check mark here if English language Translation is attached.

		NON PATENT LITERATURE DOCUMENTS	
Examiner Initials	Cite No.1	the item (book magazine journal serial symposium catalog etc.) date nage(s) volume-issue	
	CT1 ABLAMUNITS et al., Acceleration of autoimmune diabetes by cyclophosphamide is associate with an enhanced IFN-gamma secretion pathway, J. Autoimmun. 13(4):383-392 (1999)		
	CU1	J1 AHRENS et al., In vivo imaging platform for tracking immunotherapeutic cells, Nat. Biotechnol. 23(8):983-987 (2005)	
	CV1	ALLEN et al., Cellular delivery of MRI contrast agents, Chem. Bio. 11(3):301-307 (2004)	
	CW1	ANDERSON et al., Magnetic resonance imaging of labeled T-cells in a mouse model of multiple sclerosis, Ann. Neurol. 55(5):654-659 (2004)	

Complete if Known Substitute for form 1449/PTO 10/586,015 Application Number **INFORMATION DISCLOSURE** August 4, 2008 Filing Date **STATEMENT BY APPLICANT** First Named Inventor Eric T. Ahrens 1618 Art Unit (Use as many sheets as necessary) Jagadishwar Rao Samala Examiner Name 2 4 CAMU-P01-002 Sheet of Attorney Docket Number

CX1	ARBAB et al., Efficient magnetic cell labeling with protamine sulfate complexed to ferumoxides for cellular MRI, Blood 15:104(4):1217-23 (2004)	
CY1	BASSE-LUSEBRINK et al., Multi-color <sup>19</sup> F CSI: Simultaneous detection of differently labeled	
	cells in vivo, Abstract #806, Proc. Int. Soc. Mag. Reson. 17 (2009)	
CZ1	BILLOTEY et al., T-cell homing to the pancreas in autoimmune mouse models of diabetes: in vivo MR imaging, Radiology 236(2):579-587 (2005)	
CA2	BULTE et al., Preparation of magnetically labeled cells for cell tracking by magnetic resonance	
	imaging. Method Enzymol. 386:275-299 (2004)	
CB2	CANTOR et al., Effector function of diabetogenic CD4 Th1 T cell clones: a central role for TNF-alpha, J. Immunol. 175(11):7738-7745 (2005)	
CC2	CARUTHERS et al., In vitro demonstration using 19F magnetic resonance to augment	
	molecular imaging with paramagnetic perfluorocarbon nanoparticles at 1.5 Tesla, Invest.	
	Radiology 41(3):305-312	
CD2	CHENG et al., Characterization of aqueous dispersions of Fe(3)O(4) nanoparticles and their biomedical applications, Biomaterials 26(7):729-738 (2005)	
CE2	CROWDER, et al. "Unique perflourocarbon nanobeacons improve stem/progenitor cell	
	tracking with MRI" FASEB Journal, vol. 20, no. 4, part 1. March 2006, pp.A633, Abstract.	
CF2	CUNNINGHAM et al., Positive contrast magnetic resonance imaging of cells labeled with	
	magnetic nanoparticles, Mag. Res. In Med. 53:999-1005 (2005)	
CG2	ELSTER et al., Dyke Award. Europium-DTPA: a gadolinium analogue traceable by	
	fluorescence microscopy, Am. J. Neuroradiol. 10(6):1137-1144 (1989)	
CH2	EVGENOV et al., In vivo imaging of immune rejection in transplanted pancreatic islets,	
	Diabetes 55(9):2419-2428 (2006)	
Cl2	EVGENOV et al., In vivo imaging of islet transplantation, Nat. Med. 12(1):144-148 (2006)	
CJ2	FABIEN et al., Pancreatic lymph nodes are early targets of T cells during adoptive transfer of diabetes in NOD mice, J. Autoimmun. 8(3):323-334 (1995)	
CK2	FAN et al., MRI of perfluorocarbon emulsion kinetics in rodent mammary tumours, Phys. Med. & Biol. 51:211-220 (2006)	
CL2	FLÖGEL et al., In vivo monitoring of inflammation after cardiac and cerebral ischemia by	
OL2	fluorine magnetic resonance imaging, Circulation 118:140-148 (2008)	
CM2	FLORIS et al., Blood-brain barrier permeability and monocyte infiltration in experimental	
OIVIZ	allergic encephalomyelitis: a quantitative MRI study, Brain. 127(Pt 3):616-27 (2004)	
CN2	GRANOT et al., Labeling fibroblasts with biotin-BSA-GdDTPA-FAM for tracking of tumor-	
02	associated stroma by fluorescence and MR imaging, Magn, Reson. Med. 54(4):789-797 (2005)	
CO2	GUDBJARTSSON et al., The Rician distribution of noisy MRI data, Magn. Reson. Med.	
002	34(6):910-914 (1995)	
CP2	HELMER et al. On the correlation between the water diffusion coefficient and oxygen tension	
	in RIF-1 tumors, NMR in Biomedicine, 11(3):120-130 (1998)	
CQ2	HITCHENS et al., Comparison of iron-oxide- and perfluorocarbon-based cellular contrast	_
	agents for detecting immune cell infiltration in models of organ transplant rejection, Abstract	
	#931, Proc. Int. Soc. Mag. Reson. 17 (2009)	
CR2	JANJIC et al., Self-delivering nanoemulsions for dual fluorine-19 MRI and fluorescence	
	detection, J. Amer. Chem. Soc. 130:2832-2841 (2008)	
CS2		
	Fluorinated Oils and Amphiles, Tetrahedron 63(19):3982-3988 (2007)	
CT2	KIM et al., Interplay of tumor vascular oxygenation and tumor pO2 observed using near-	
	infrared spectroscopy, an oxygen needle electrode, and 19F MR pO2 mapping, J. Biomed Opt 8:53-62; 2003	
CU2	KIRCHER et al., In vivo high resolution three-dimensional imaging of antigen-specific cytotoxic	
	T-lymphocyte trafficking to tumors, Cancer Res. 63(20):6838-6846 (2003)	
CV2	KLUG et al., 1H/19F molecular MR-imaging in mouse models of acute and chronic	
	inflammation, Abstract #3172, Proc. Int. Soc. Mag. Reson. 17 (2009)	

Complete if Known Substitute for form 1449/PTO Application Number 10/586,015 **INFORMATION DISCLOSURE** August 4, 2008 Filing Date STATEMENT BY APPLICANT First Named Inventor Eric T. Ahrens 1618 Art Unit (Use as many sheets as necessary) Jagadishwar Rao Samala Examiner Name 3 4 CAMU-P01-002 Sheet of Attorney Docket Number

CW2 KRAITCHMAN et al., In vivo magnetic resonance imaging of mesenchymal stem cells in myocardial infarction, Circulation 107(18);2290-2293 (2003)  CX2 KRAVTZOFF et al., GD-DOTA Loaded into red blood cells. A new magnetic resonance imaging contrast agents for vascular system," Adv. in Exp. Med. and Biol. 326:347-326 (1992)  CY2 LANZA et al., 1H/19F magnetic resonance molecular imaging with perfluorocarbon nanoparticles. In: Ahrens ET, editor. In vivo cellular and molecular imaging, Curr. Top. Dev. Biol. 70:58-78 (2005)  CX2 LANZA et al., A novel site-targeted ultrasonic contrast agent with broad biomedical application, Circulation 34(12):3334-3340 (1996)  CA3 LAUKEMPER-OSTENDORF et al., 19F-MRI of perfluoror for measurement of oxygen partial pressure in porcine lungs during partial liquid ventilation, Magn. Reson. Med. 47:82-89; 2002  CB3 LEITER et al., The nonobese diabetic (NOD) mouse, Am. J. Pathol. 128(2):380-383 (1987)  CC3 MASON et al., Hexafluorobenzene: a sensitive 19F NMR indicator of tumor oxygenation, NMR Biomed 9:125-134; 1996  CD3 McNAB et al., Tissue oxygen tension measurements in the Shionogi model of prostate cancer using "F MRS and MRI, MAGMA 17:288-295 (2004)  CE3 MEYER et al., Measurement of vascular volume in experimental rat tumors by 19F magnetic resonance imaging, Invest. Radiol. 28(6);710-719 (1993)  CF3 MILLER et al., Imaging the single cell dynamics of CD4+ T cell activation by dendritic cells in lymph nodes, J. Exp. Med. 200(7):847-856 (2004)  GA3 MIYAZAKI et al., Predominance of lymphocytes-1 in pancreatic-islets and spleen of prediabetic non-obese diabetic (NOD) mice - a longitudinal-study, Clin. Exp. Immunol. 60(3):622-630 (1985)  GH3 MODO et al., Mapping transplanted stem cell migration after a stroke; a serial, in vivo magnetic resonance imaging study, Neuroimage 21(1):311-317 (2004)  GN3 MODO et al., Tracking the recruitment of diabetogenic CP8+ T-cells to the pancreas in real time, Diabetes 53(6):1459-1466 (2004)  GN3 MODO et al., Mapping transplanted stem cell migration after		
CX2 KRAVTZOFF et al., GD-DOTA Loaded into red blood cells. A new magnetic resonance imaging contrast agents for vascular system," Adv. in Exp. Med. and Biol. 326:347-326 (1992)  CY2 LANZA et al., 1H/19F magnetic resonance molecular imaging with perfluorocarbon nanoparticles. In: Ahrens ET, editor. In vivo cellular and molecular imaging, Curr. Top. Dev. Biol. 70:88-78 (2005)  CZ2 LANZA et al., A novel site-targeted ultrasonic contrast agent with broad biomedical application, Circulation py4(12):3334-3340 (1996)  CA3 LAUKEMPER-OSTENDORF et al., 19F-MRI of perflubron for measurement of xygen partial pressure in porcine lungs during partial liguid venilitation, Magn. Reson. Med. 47:82-89; 2002  CB3 LEITER et al., The nonobese diabetic (NOD) mouse, Am. J. Pathol. 128(2):380-383 (1997)  CC3 MASON et al., Hexafluorobenzene: a sensitive 19F NMR indicator of tumor oxygenation, NMR Biomed 9:125-134; 1996  MeNAB et al., Tissue oxygen lension measurements in the Shionogi model of prostate cancer using "F MRS and MRI, MAGMA 17:288-295 (2004)  CE3 MEYER et al., Measurement of vascular volume in experimental rat tumors by 19F magnetic resonance imaging, Invest. Radiol. 28(8):710-719 (1993)  CF3 MILLER et al., Imaging the single cell dynamics of CD4+ T cell activation by dendritic cells in lymph nodes. J. Exp. Med. 2007(3:847-856 (2004)  CG3 MYAZAKI et al., Predominance of lymphocytes-T in pancreatic-slets and spleen of prediabetic non-obese diabetic (NOD) mice – a longitudinal-study, Clin. Exp. Immunol. 60(3):622-630 (1985)  CH3 MOO et al., Mapping transplanted stem cell migration after a stroke: a serial, in vivo magnetic resonance imaging study, Neuroimage 21(1):311-317 (2004)  MOO et al., Quantitative magnetic resonance immunohistochemistry with ligand-targeted F-19 nanoparticles. Magn. Reson. Med. 52(6):1255-1262 (2004)  MOORE et al., Tracking the recruitment of diabetogenic CD8+ T-cells to the pancreasi in real time, Diabetes 53(6):1459-1466 (2004)  MOORE et al., Trapeted Nanoparticles for Quantitative maging of Sparse M	CW:	
imaging contrast agents for vascular system," Adv. in Exp. Med. and Biol. 326:347-326 (1992)  CY2 LANZA et al., 1H/19F magnetic resonance molecular imaging with perfluorocarbon nanoparticles. In: Ahrens ET, editor. In vivo cellular and molecular imaging, Curr. Top. Dev. Biol. 70:58-78 (2005)  CZ2 LANZA et al., A novel site-targeted ultrasonic contrast agent with broad biomedical application, Circulation 94(12):3334-3340 (1996)  CA3 LAUKEMPER-OSTENDORT et al., 19F-MRI of perflubron for measurement of oxygen partial pressure in porcine lungs during partial liquid ventilation, Magn. Reson. Med. 47:82-89; 2002  CB3 LETTER et al., The nonbose diabetic (NOD) mous. Am. J. Pathol. 128(2):380-3383 (1987)  CC3 MASON et al., Hexafluorobenzene: a sensitive 19F NMR indicator of tumor oxygenation, NMR Biomed 9:125-134; 1996  MNAB et al., Tissue oxygen tension measurements in the Shionogi model of prostate cancer using "F MRS and MRI, MAGMA 17:288-295 (2004)  GE3 MEYER et al., Measurement of vascular volume in experimental rat tumors by 19F magnetic resonance imaging, Invest. Radiol. 28(8):710-719 (1993)  GF3 MILLER et al., Imaging the single cell drynamics of CO4+ T cell activation by dendritic cells in lymph nodes, J. Exp. Med. 200(7):847-856 (2004)  MYAZAKI et al., Fredominance of lymphocytes-T in pancreatic-islets and spleen of prediabetic non-obese diabetic (NOD) mice – a longitudinal-study, Clin. Exp. Immunol. 60(3):622-630 (1985)  CH3 MODG et al., Mapping transplanted stem cell migration after a stroke: a serial, in vivo magnetic resonance imaging study, Neuroimage 21(1):311-317 (2004)  MOARSH et al., Tracking the recruitment of diabetogenic CD8+ T-cells to the pancreas in real time, Diabetes 53(6):1459-1466 (2004)  MOARSH et al., Trangeted Nanoparticles for Quantitative maging of Sparse Molecular Epitopes with MRI, Mag. Res. in Med. 51(3):840-848 (2004)  CX3 MORAWSKI et al., Targeted Nanoparticles for Quantitative maging of Sparse Molecular Epitopes with MRI, Mag. Res. in Med. 51(3):480-948 (2004)  CX3 MORAWSKI et al.	CV	
nanoparticles. In: Ahrens ET, editor. In vivo cellular and molecular imaging, Curr. Top. Dev. Biol. 70:58-78 (2005)  C22 LANZA et al., A novel site-targeted ultrasonic contrast agent with broad biomedical application, Circulation 94(12):3334-3340 (1996)  CA3 LAUKEMPER-OSTENDOFF et al., 199-MRI of perflubron for measurement of oxygen partial pressure in porcine lungs during partial liquid ventilation, Magn. Reson. Med. 47:82-89; 2002  CB3 LETIER et al., The nonobese diabetic (NDD) mouse, Am. J. Pathol. 128(2):380-383 (1997)  CC3 MASON et al., Hexafluorobenzene: a sensitive 19F NMR indicator of tumor oxygenation, NMR Biomed 9:125-134; 1996  CD3 McNAB et al., Tissue oxygen tension measurements in the Shionogi model of prostate cancer using "F MRS and MRI, MAGMA 17:288-295 (2004)  CC3 MEYER et al., Measurement of vascular volume in experimental rat tumors by 19F magnetic resonance imaging, Invest. Radiol. 28(8):710-719 (1993)  GF3 MILLER et al., Imaging the single cell drynamics of CD4 + T cell activation by dendritic cells in lymph nodes, J. Exp. Med. 200(7):847-856 (2004)  GG3 MIYAZAKI et al., Predominance of lymphocytes-T in pancreatic-islets and spleen of prediabetic non-obese diabetic (NCD) mice - a longitudinal-study, Clin. Exp. Immunol. 60(3):622-630 (1985)  CH3 MODO et al., Mapping transplanted stem cell migration after a stroke: a serial, in vivo magnetic resonance imaging study, Neuroimage 21(1):311-317 (2004)  MOORE et al., Tracking the recruitment of diabetogenic CD8+ T-cells to the pancreas in real time, Diabetes 53(6):1459-1466 (2004)  MORAWSKI et al., Quantitative magnetic resonance immunohistochemistry with ligand-largeted F-19 nanoparticles, Magn. Reson. Med. 52(6):1255-1262 (2004)  MORAWSKI et al., Endothelial stem cell detection in vivo with unique perfluorocarbon nanoparticle labels using fluorine (F-19) MNRI at 1.5 T, Circulation 114(18)(Suppl. S): 251 (Abstract) (2006)  PAKTLOW et al., 19F magnetic resonance imaging for stemplrogenitor cell tracking with multiple unique perfluorocarbon nanobea		imaging contrast agents for vascular system," Adv. in Exp. Med. and Biol. 326:347-326 (1992)
Biol. 70:58-78 (2005)  C22 LANZA et al., A novel site-targeted ultrasonic contrast agent with broad biomedical application, Circulation 94(12):3334-3340 (1996)  C33 LAUKEMPER-OSTENDORF et al., 19F-MRI of perflubron for measurement of oxygen partial pressure in portine lungs during partial liquid ventilation, Magn. Reson. Med. 47:82-99; 2002  C83 LEITER et al., The nonobese diabetic (NOD) mouse, Am. J. Pathol. 128(2):380-383 (1987)  CC3 MASON et al., Hexafluorobenzene: a sensitive 19F NMR indicator of tumor oxygenation, NMR Biomed 9:125-134; 1996  McNAB et al., Tissue oxygen tension measurements in the Shionogi model of prostate cancer using "F MRS and MRI, MAGMA 17:288-295 (2004)  C83 MEYER et al., Measurement of vascular volume in experimental rat tumors by 19F magnetic resonance imaging, Invest. Radiol. 28(8):710-719 (1993)  C73 MILLER et al., Imaging the single cell dynamics of CD4+ T cell activation by dendritic cells in lymph nodes, J. Exp. Med. 200(7):847-856 (2004)  C33 MIYAZAKI et al., Predominance of lymphocytes-T in pancreatic-islets and spleen of prediabetic non-obese diabetic (NOD) mice - a longitudinal-study, Clin. Exp. Immunol. 60(3):622-630 (1985)  C43 MODO et al., Mapping transplanted stem cell migration after a stroke: a serial, in vivo magnetic resonance imaging, study, Neuroimage 21(1):311-317 (2004)  C33 MORAWSKI et al., Tracking the recruitment of diabetogenic CD8+ T-cells to the pancreas in real time, Diabetes 53 (6):1459-1466 (2004)  C43 MORAWSKI et al., Targeted Annoparticles for Quantitative magning of Sparse Molecular Epitopes with MRK, Mag. Res. in Med. 51(3):480-486 (2004)  C43 MORAWSKI et al., Targeted Inanoparticles for Quantitative magning of Sparse Molecular Epitopes with MRK, Mag. Res. in Med. 51(3):480-486 (2004)  C53 MORAWSKI et al., Targeted nanoparticles for Quantitative magning of Sparse Molecular Epitopes with MRK, Mag. Res. in Med. 51(3):480-486 (2004)  C63 PiACLA et al., The pragnetic resonance imaging for stem/progenior cell tracking with multiple unique perfluorocarbon	CY2	
C22 LANZA et al., A novel site-targeted ultrasonic contrast agent with broad biomedical application, Circulation 94(12):3334-3340 (1996)  CA3 LAUKEMPER-OSTENDORF et al., 19F-MRI of perflubron for measurement of oxygen partial pressure in porcine lungs during partial liquid ventilation, Magn. Reson. Med. 47:82-99; 2002  CB3 LEITER et al., The nonobese diabetic (NOD) mouse, Am. J. Pathol. 128(2):380-383 (1987)  CC3 MASON et al., Hexafluorobenzene: a sensitive 19F NMR indicator of tumor oxygenation, NMR Biomed 9:125-134; 1996  CD3 McNAB et al., Tissue oxygen tension measurements in the Shionogi model of prostate cancer using <sup>18</sup> F MRS and MRI, MAGMA 17:288-295 (2004)  CE3 MEYER et al., Measurement of vascular volume in experimental rat tumors by 19F magnetic resonance imaging, Invest. Radiol. 28(8):710-719 (1993)  CF3 MILLER et al., Imaging the single cell dynamics of CD4+ T cell activation by dendritic cells in lymph nodes, J. Exp. Med. 200(7):847-856 (2004)  CG3 MIYAZAKI et al., Predominance of lymphocytes-T in pancreatic-islets and spleen of prediabetic non-obese diabetic (NOD) mice – a longitudinal-study, Clin. Exp. Immunol. 60(3):622-630 (1985)  CH3 MODO et al., Mapping transplanted stem cell migration after a stroke: a serial, in vivo magnetic resonance imaging study, Neuroimage 21(1):311-317 (2004)  G13 MODRE et al., Tracking the recruitment of diabetogenic CD8+ T-cells to the pancreas in real time, Diabetes 53(6):1459-1466 (2004)  CK3 MORAWSKI et al., Quantitative magnetic resonance immunohistochemistry with ligand-targeted F-19 nanoparticles, Magn. Reson. Med. 52(6):1255-1262 (2004)  CK3 MORAWSKI et al., Targeted Nanoparticles for Quantitative Imaging of Sparse Molecular Epitopes with MRK, Mag. Res. in Med. 51(3):480-486 (2004)  CK3 PAKALA et al., Thelper 2 (Th2) T cells induce acute pancreatitis and diabetes in immune-compromised nonobese diabetic (NOD) mice, J. Exp. Med. 186(2):299-306 (1997)  CN3 PARTLOW et al., 19F magnetic resonance imaging for stem/progenitor cell tracking with multiple unique perfl		
Circulation 94(12):3334-3340 (1996)  CA3 LAUKEMPER-OSTENDORF et al., 19F-MRI of perflubron for measurement of oxygen partial pressure in porcine lungs during partial liquid ventilation, Magn. Reson. Med. 47:82-89; 2002  CB3 LEITER et al., The nonobese diabetic (NOD) mouse, Am. J. Pathol. 128(2):380-383 (1987)  CC3 MASON et al., Hexafluorobenzene: a sensitive 19F MMR indicator of tumor oxygenation, NMR Biomed 9:125-134; 1996  CD3 MCNAB et al., Tissue oxygen tension measurements in the Shionogi model of prostate cancer using <sup>19</sup> F MRS and MRI, MAGMA 17:288-295 (2004)  CE3 MEYER et al., Measurement of vascular volume in experimental rat tumors by 19F magnetic resonance imaging, Invest. Radiol. 26(8):710-719 (1993)  CF3 MILLER et al., Imaging the single cell dynamics of CD4+ T cell activation by dendritic cells in lymph nodes, J. Exp. Med. 2007(19:47-356 (2004)  CG3 MIYAZAKI et al., Predominance of lymphocytes-T in pancreatic-slets and spleen of prediabetic non-obese diabetic (NOD) mice – a longitudinal-study. Clin. Exp. Immunol. 60(3):622-630 (1985)  CH3 MODO et al. Mapping transplanted stem cell migration after a stroke: a serial, in vivo magnetic resonance imaging study, Neuroimage 21(1):311-317 (2004)  CI3 MORAWSKI et al., Tracking the recruitment of diabetogenic CD8+ T-cells to the pancreas in real time. Diabetes 53(6):1459-1466 (2004)  CK3 MORAWSKI et al., Targeted Nanoparticles for Quantitative Imaging of Sparse Molecular Epitopes with MRK, Mag. Res. in Med. 51(3):480-486 (2004)  CK3 MORAWSKI et al., Targeted Nanoparticles for Quantitative Imaging of Sparse Molecular Epitopes with MRK, Mag. Res. in Med. 51(3):480-486 (2004)  CL3 NEURALTER et al., Endothelial stem cell detection in vivo with unique perfluorocarbon nanoparticle labets using fluorine (F-19) MNRI at 1.5 T, Circulation 114(18)(Suppl. S): 251 (Abstract) (2006)  CM3 PAKALA et al., Thelper 2 (Th2) T cells induce acute pancreatitis and diabetes in immunecompromised nonobese diabetic (NOD) mice, J. Exp. Med. 186(2):299-306 (1997)  CN3 PARTLOW et al.,		
CA3 LAUKEMPER-OSTENDORF et al., 19F-NRI of perflutron for measurement of oxygen partial pressure in porcine lungs during partial liquid ventilation, Magn. Reson. Med. 47:82-89; 2002  CB3 LEITER et al., The nonobese diabetic (NOD) mouse, Am. J. Pathol. 128(2):380-383 (1987)  CC3 MASON et al., Hexafluorobenzene: a sensitive 19F NMR indicator of tumor oxygenation, NMR Biomed 9:125-134; 1996  CD3 McNAB et al., Trissue oxygen tension measurements in the Shionogi model of prostate cancer using <sup>19</sup> F MRS and MRI, MAGMA 17:288-295 (2004)  CE3 MEYER et al., Measurement of vascular volume in experimental rat tumors by 19F magnetic resonance imaging, Invest. Radiol. 28(8):710-719 (1993)  CF3 MILLER et al., ingrigh the single cell dynamics of CD4+ T cell activation by dendritic cells in lymph nodes. J. Exp. Med. 200(7):847-856 (2004)  CG3 MIYAZAKI et al., Predominance of lymphocytes-T in pancreatic-islets and spleen of prediabetic non-obese diabetic (NOD) mice - a longitudinal-study, Clin. Exp. Immunol. 60(3):622-630 (1985)  CH3 MODO et al., Mapping transplanted stem cell migration after a stroke: a serial, in vivo magnetic resonance imaging study, Neuroimage 21(1):311-317 (2004)  CI3 MORR et al., Tracking the recruitment of diabetogenic CD8+ T-cells to the pancreas in real time, Diabetes 53(6):1459-1466 (2004)  CJ3 MORAWSKI et al., Quantitative magnetic resonance immunohistochemistry with ligand-targeted F-19 nanoparticles. Magn. Reson. Med. 52(6):1255-1262 (2004)  CK3 MORAWSKI et al., Targeted Nanoparticles for Cuantitative Imaging of Sparse Molecular Epitopes with MRK, Mag. Res. in Med. 51(3):480-486 (2004)  CK3 MORAWSKI et al., Trageted Nanoparticles for Cuantitative Imaging of Sparse Molecular Epitopes with MRK, Mag. Res. in Med. 51(3):480-486 (2004)  CK3 PAKALA et al., Thelper 2 (Th2) T cells induce acute pancrealitis and diabetes in immune-compromised nonobese diabetic (NOD) mice, J. Exp. Med. 1786(2):299-306 (1997)  CM3 PARTLOW et al., 19F magnetic resonance imaging for stem/progenitor cell tracking with multiple u	CZ2	
pressure in porcine lungs during partial liquid ventilation, Magn. Reson. Med. 47:82-89; 2002  CB3 LEITER et al., The nonobese diabetic (NOD) mouse, Am. J. Pathol. 128(2):380-383 (1987)  CC3 MASON et al., Hexafluorobenzene: a sensitive 19F MMR indicator of tumor oxygenation, NMR Biomed 9:125-134; 1996  CD3 MCNAB et al., Tissue oxygen tension measurements in the Shionogi model of prostate cancer using <sup>19</sup> F MRS and MRI, MAGMA 17:288-295 (2004)  CB3 MEYER et al., Measurement of vascular volume in experimental rat tumors by 19F magnetic resonance imaging, Invest. Radiol. 28(8):710-719 (1993)  CF3 MILLER et al., imaging the single cell dynamics of CD4+ T cell activation by dendritic cells in lymph nodes. J. Exp. Med. 200(7):847-856 (2004)  CG3 MIYAZAKI et al., Predominance of lymphocytes-T in pancreatic-islets and spleen of prediabetic non-obese diabetic (NOD) mice – a longitudinal-study, Clin. Exp. Immunol. 60(3):622-630 (1985)  CH3 MODO et al., Mapping transplanted stem cell migration after a stroke: a serial, in vivo magnetic resonance imaging study, Neuroimage 21(1):311-317 (2004)  CI3 MORE et al., Tracking the recruitment of diabetogenic CD8+ T-cells to the pancreas in real time, Diabetes 53(6):1459-1466 (2004)  CJ3 MORAWSKI et al., Quantitative magnetic resonance immunohistochemistry with ligand-targeted F-19 nanoparticles, Magn. Reson. Med. 52(6):1255-1262 (2004)  CK3 MORAWSKI et al., Cuantitative magnetic resonance immunohistochemistry with ligand-targeted F-19 nanoparticles in Magn. Reson. Med. 51(3):480-486 (2004)  CL3 REUBAUER et al., Endothelial stem cell detection in vivo with unique perfluorocarbon nanoparticle labels using fluorine (F-19) MNRI at 1.5 T, Circulation 114(18)(Suppl. S): 251 (Abstract) (2006)  CM3 PAKALA et al., Thelper 2 (Th2) T cells induce acute pancreatitis and diabetes in immune-compromised nonobese diabetic (NOD) mice, J. Exp. Med. 186(2):299-306 (1997)  CN3 PARTLOW et al., 19F magnetic resonance imaging for stem/progenitor cell tracking with multiple unique perfluorocarbon nanobea	CA3	
CC3 MASON et al., Hexafluorobenzene: a sensitive 19F NMR indicator of tumor oxygenation, NMR Biomed 9:125-134; 1996  CD3 MCNAB et al., Tissue oxygen tension measurements in the Shionogi model of prostate cancer using <sup>16</sup> F MRS and MRI, MAGMA 17:288-295 (2004)  CE3 MEYER et al., Measurement of vascular volume in experimental rat tumors by 19F magnetic resonance imaging, Invest. Radiol. 28(8):710-719 (1993)  CF3 MILLER et al., Imaging the single cell dynamics of CD4+ T cell activation by dendritic cells in lymph nodes, J. Exp. Med. 200(7):847-856 (2004)  CG3 MIYAZAKI et al., Predominance of lymphocytes-T in pancreatic-islets and spleen of prediabetic non-obese diabetic (NOD) mice - a longitudinal-study, Clin. Exp. Immunol. 60(3):622-630 (1985)  CH3 MODO et al., Mapping transplanted stem cell migration after a stroke: a serial, in vivo magnetic resonance imaging study, Neuroimage 21(1):311-317 (2004)  CI3 MORAWSKI et al., Cuantitative magnetic resonance immunohistochemistry with ligand-targeted F-19 nanoparticles, Magn. Reson. Med. 52(6):1255-1262 (2004)  CK3 MORAWSKI et al., Cuantitative magnetic resonance immunohistochemistry with ligand-targeted F-19 nanoparticles, Magn. Reson. Med. 52(6):1255-1262 (2004)  CK3 MORAWSKI et al., Endothelial stem cell detection in vivo with unique perflourocarbon nanoparticle labels using fluorine (F-19) MNRI at 1.5 T, Circulation 114(18)(Suppl. S): 251 (Abstract) (2006)  CM3 PAKALA et al., Thelper 2 (Th2) T cells induce acute pancreatitis and diabetes in immune-compromised nonobese diabetic (NOD) mice, J. Exp. Med. 186(2):299-306 (1997)  CN3 PARTLOW et al., 19F magnetic resonance imaging for stem/progenitor cell tracking with multiple unique perfluorocarbon nanobarcoting of stem-progenitor cell tracking with multiple unique perfluorocarbon nanobaecoms. FASEB J. 21:1647-1654 (2007)  PELCHEN-MATTHEWS et al., Phorbol ester-induced downregulation of CD4 is a multistep process involving dissociation from p56ick, increased association with clathrin-coated pits, and altered endosomal so		
Biomed 9:125-134; 1996  CD3 McNAB et al., Tissue oxygen tension measurements in the Shionogi model of prostate cancer using "F MRS and MRI, MAGMA 17:288-295 (2004)  CE3 MEYER et al., Measurement of vascular volume in experimental rat tumors by 19F magnetic resonance imaging, Invest. Radiol. 28(8):710-719 (1993)  CF3 MILLER et al., Imaging the single cell dynamics of CD4+ T cell activation by dendritic cells in lymph nodes, J. Exp. Med. 200(7):847-856 (2004)  CG3 MIYAZAKI et al., Predominance of lymphocytes-T in pancreatic-islets and spleen of prediabetic non-obese diabetic (NOD) mice - a longitudinal-study, Clin. Exp. Immunol. 60(3):622-630 (1985)  CH3 MODO et al., Mapping transplanted stem cell migration after a stroke: a serial, in vivo magnetic resonance imaging study. Neuroimage 21(1):311-317 (2004)  CI3 MORAWSKI et al., Tracking the recruitment of diabetogenic CD8+ T-cells to the pancreas in real time, Diabetes 53(6):1459-1466 (2004)  CJ3 MORAWSKI et al., Quantitative magnetic resonance immunohistochemistry with ligand-targeted F-19 nanoparticles, Magn. Reson. Med. 52(6):1255-1262 (2004)  CK3 MORAWSKI et al., Targeted Nanoparticles for Quantitative Imaging of Sparse Molecular Epitopes with MRK, Mag. Res. in Med. 51(3):480-486 (2004)  CL3 MORAWSKI et al., Targeted Nanoparticles for Quantitative Imaging of Sparse Molecular Epitopes with MRK, Mag. Res. in Med. 51(3):480-486 (2004)  CL3 MORAWSKI et al., Tandothelial stem cell detection in vivo with unique perflourocarbon nanoparticle labels using fluorine (F-19) MNRI at 1.5 T, Circulation 114(18)(Suppl. S): 251 (Abstract) (2006)  CM3 PAKALA et al., Thelper 2 (Th2) T cells induce acute pancreatitis and diabetes in immunecompromised nonobese diabetic (NOD) mice, J. Exp. Med. 186(2):299-306 (1997)  PARTLOW et al., 19F magnetic resonance imaging for stem/propenitor cell tracking with multiple unique perfluorocarbon nanobeacons. FASEB J. 21:1647-1654 (2007)  PARTLLIPS et al., Nondepleting anti-CD4 has an immediate action on diabetogenic effector cells, halting thei	CB3	LEITER et al., The nonobese diabetic (NOD) mouse, Am. J. Pathol. 128(2):380-383 (1987)
CD3 MCNAB et al., Tissue oxygen tension measurements in the Shionogi model of prostate cancer using <sup>19</sup> F MRS and MRI, MAGMA 17:288-295 (2004)  CE3 MEYER et al., Measurement of vascular volume in experimental rat tumors by 19F magnetic resonance imaging, Invest. Radiol. 28(8):710-719 (1993)  CF3 MILLER et al., Imaging the single cell dynamics of CD4+ T cell activation by dendritic cells in lymph nodes, J. Exp. Med. 200(7):847-856 (2004)  CG3 MIYAZAKI et al., Predominance of lymphocytes-T in pancreatic-islets and spleen of prediabetic non-obese diabetic (NOD) mice – a longitudinal-study, Clin. Exp. Immunol. 60(3):622-630 (1985)  CH3 MODO et al., Mapping transplanted stem cell migration after a stroke: a serial, in vivo magnetic resonance imaging study, Neuroimage 21(1):311-317 (2004)  MODORE et al., Tracking the recruitment of diabetogenic CD8+ T-cells to the pancreas in real time, Diabetes 53(6):1459-1466 (2004)  CJ3 MORAWSKI et al., Quantitative magnetic resonance immunohistochemistry with ligand-targeted F-19 nanoparticles, Magn. Reson. Med. 52(6):1255-1262 (2004)  CK3 MORAWSKI et al., Targeted Nanoparticles for Quantitative Imaging of Sparse Molecular Epitopes with MRK, Mag. Res. in Med. 51(3):480-486 (2004)  CL3 MORAWSKI et al., Endothelial stem cell detection in vivo with unique perflourocarbon nanoparticle labels using fluorine (F-19) MNRI at 1.5 T, Circulation 114(18)(Suppl. S): 251 (Abstract) (2006)  CM3 PAKALA et al., T helper 2 (Th2) T cells induce acute pancreatitis and diabetes in immune-compromised nonobese diabetic (NOD) mice, J. Exp. Med. 186(2):299-306 (1997)  CN3 PARTLOW et al., 19F magnetic resonance imaging for stemplrogenitor cell tracking with multiple unique perfluorocarbon nanobeacons, FASEB J. 21:1647-1654 (2007)  CO3 PELCHEN-MATTHEWS et al., Phorbol ester-induced downregulation of CD4 is a multistep process involving dissociation from p56lck, increased association with clathrin-coated pits, and altered endosomal sorting, J. Exp. Med. 178(4):1209-1222 (1993)  PHILLIPS et al., Nondepletin	CCS	MASON et al., Hexafluorobenzene: a sensitive 19F NMR indicator of tumor oxygenation, NMR
using <sup>18</sup> F MRS and MRI, MAGMA 17:288-295 (2004)  CE3 MEYER et al., Measurement of vascular volume in experimental rat tumors by 19F magnetic resonance imaging, Invest. Radiol. 28(8):710-719 (1993)  CF3 MILLER et al., Imaging the single cell dynamics of CD4+ T cell activation by dendritic cells in lymph nodes, J. Exp. Med. 200(7):847-856 (2004)  CG3 MIYAZAKI et al., Predominance of lymphocytes-T in pancreatic-islets and spleen of prediabetic non-obese diabetic (NOD) mice - a longitudinal-study, Clin. Exp. Immunol. 60(3):622-630 (1985)  CH3 MODO et al., Mapping transplanted stem cell migration after a stroke: a serial, in vivo magnetic resonance imaging study, Neuroimage 21(1):311-317 (2004)  CI3 MODRA et al., Tracking the recruitment of diabetogenic CD8+ T-cells to the pancreas in real time, Diabetes 53(6):1459-1466 (2004)  CJ3 MORAWSKI et al., Quantitative magnetic resonance immunonistochemistry with ligand-targeted F-19 nanoparticles, Magn. Reson. Med. 52(6):1255-1262 (2004)  CK3 MORAWSKI et al., Targeted Nanoparticles for Quantitative Imaging of Sparse Molecular Epitopes with MRK, Mag. Res. in Med. 51(3):480-486 (2004)  CL3 NEUBAUER et al., Targeted Nanoparticles for Quantitative Imaging of Sparse Molecular Epitopes with MRK, Mag. Res. in Med. 51(3):480-486 (2004)  CL3 NEUBAUER et al., Endothelial stem cell detection in vivo with unique perflourocarbon nanoparticle labels using fluorine (F-19) MNRI at 1.5 T, Circulation 114(18)(Suppl. S): 251 (Abstract) (2006)  CM3 PAKALA et al., Thelper 2 (Th2) T cells induce acute pancreatitis and diabetes in immune-compromised nonobese diabetic (NOD) mice, J. Exp. Med. 186(2):299-306 (1997)  CN3 PARTLOW et al., 19F magnetic resonance imaging for stem/progenitor cell tracking with multiple unique perfluorocarbon nanobeacons, FASEB J. 21:1647-1654 (2007)  CO3 PELCHEN-MATHEWS et al., Phorbol ester-induced downrepation of CD4 is a multistep process involving dissociation from p56l6k, increased association with clathrin-coated pits, and altered endosomal sorting, J. Exp. Med.		
CE3 MEYER et al., Measurement of vascular volume in experimental rat tumors by 19F magnetic resonance imaging, Invest. Radiol. 28(8):710-719 (1993)  CF3 MILLER et al., Imaging the single cell dynamics of CD4+T cell activation by dendritic cells in lymph nodes. J. Exp. Med. 200(7):847-856 (2004)  CG3 MIYAZAKI et al., Predominance of lymphocytes-T in pancreatic-islets and spleen of prediabetic non-obese diabetic (NOD) mice - a longitudinal-study, Clin. Exp. Immunol. 60(3):622-630 (1985)  CH3 MODO et al., Mapping transplanted stem cell migration after a stroke: a serial, in vivo magnetic resonance imaging study, Neuroimage 21(1):311-317 (2004)  CI3 MOORE et al., Tracking the recruitment of diabetogenic CD8+ T-cells to the pancreas in real time, Diabetes 53(6):1459-1466 (2004)  CJ3 MORAWSKI et al., Quantitative magnetic resonance immunohistochemistry with ligand-targeted F-19 nanoparticles, Magn. Reson. Med. 52(6):1255-1262 (2004)  CK3 MORAWSKI et al., Targeted Nanoparticles for Quantitative Imaging of Sparse Molecular Epitopes with MRK, Mag. Res. in Med. 51(3):480-486 (2004)  CL3 NEUBAUER et al., Endothelial stem cell detection in vivo with unique perflourocarbon nanoparticle labels using fluorine (F-19) MNRI at 1.5 T, Circulation 114(18)(Suppl. S): 251 (Abstract) (2006)  CM3 PAKALA et al., Thelper 2 (Th2) T cells induce acute pancreatitis and diabetes in immune-compromised nonobese diabetic (NOD) mice, J. Exp. Med. 186(2):299-306 (1997)  CN3 PARTLOW et al., 19F magnetic resonance imaging for stem/progenitor cell tracking with multiple unique perfluorocarbon nanobeacons, FASEB J. 21:1647-1654 (2007)  PARTLOW et al., 19F magnetic resonance imaging for stem/progenitor cell tracking with multiple unique perfluorocarbon nanobeacons, FASEB J. 21:1647-1654 (2007)  PECHEN-MATTHEWS et al., Phorbol ester-induced downregulation of CD4 is a multistep process involving dissociation from p56lck, increased association with clathrin-coated pits, and altered endosomal sorting, J. Exp. Med. 178(4):1209-1222 (1993)  PHILLIPS et al.	CD3	MCNAB et al., Tissue oxygen tension measurements in the Shionogi model of prostate cancer
resonance imaging, Invest. Radiol. 28(8):710-719 (1993)  CF3 MILLER et al., Imaging the single cell dynamics of CD4+ T cell activation by dendritic cells in lymph nodes, J. Exp. Med. 2007):847-856 (2004)  CG3 MIYAZAKI et al., Predominance of lymphocytes-T in pancreatic-islets and spleen of prediabetic non-obese diabetic (NOD) mice - a longitudinal-study, Clin. Exp. Immunol. 60(3):622-630 (1985)  CH3 MODO et al., Mapping transplanted stem cell migration after a stroke: a serial, in vivo magnetic resonance imaging study, Neuroimage 21(1):311-317 (2004)  CI3 MODRE et al., Tracking the recruitment of diabetogenic CD8+ T-cells to the pancreas in real time, Diabetes 53(6):1459-1466 (2004)  CJ3 MORAWSKI et al., Quantitative magnetic resonance immunohistochemistry with ligand-targeted F-19 nanoparticles, Magn. Reson. Med. 52(6):1255-1262 (2004)  CK3 MORAWSKI et al., Targeted Nanoparticles for Quantitative Imaging of Sparse Molecular Epitopes with MRK, Mag. Res. in Med. 51(3):480-486 (2004)  CL3 NEUBAUER et al., Endothelial stem cell detection in vivo with unique perflourocarbon nanoparticle labels using fluorine (F-19) MNRI at 1.5 T, Circulation 114(18)(Suppl. S): 251 (Abstract) (2006)  CM3 PAKALA et al., T helper 2 (Th2) T cells induce acute pancreatitis and diabetes in immune-compromised nonobese diabetic (NOD) mice, J. Exp. Med. 186(2):299-306 (1997)  CN3 PARTLOW et al., 19F magnetic resonance imaging for stem/progenitor cell tracking with multiple unique perfluorocarbon nanobeacons, FASEB J. 21:1647-1654 (2007)  CO3 PELCHEN-MATTHEWS et al., Phorbol ester-induced downregulation of CD4 is a multistep process involving dissociation from p56ick, increased association with clathrin-coated pits, and altered endosomal sorting, J. Exp. Med. 178(4):1209-1222 (1993)  CP3 PHILLIPS et al., MAdCAM-1 is needed for diabetes development mediated by the T cell clone, BDC-2.5, Immunology 116(4):525-531 (2005)  CR3 PACENTI et al., Synthesis and characterization of fluorinated polyetheric amides, J. Fluor. Chem. 68:227-235 (1994)  CS3		USING F WIRS AND WIRI, WAGMA 17:200-295 (2004)
Iymph nodes, J. Exp. Med. 200(7):847-856 (2004)	CES	resonance imaging, Invest. Radiol. 28(8):710-719 (1993)
CG3 MIYAZAKI et al., Predominance of lymphocytes-T in pancreatic-islets and spleen of prediabetic non-obese diabetic (NOD) mice - a longitudinal-study, Clin. Exp. Immunol. 60(3):622-630 (1985)  CH3 MODO et al., Mapping transplanted stem cell migration after a stroke: a serial, in vivo magnetic resonance imaging study, Neuroimage 21(1):311-317 (2004)  CI3 MOORE et al., Tracking the recruitment of diabetogenic CD8+ T-cells to the pancreas in real time, Diabetes 53(6):1459-1466 (2004)  CJ3 MORAWSKI et al., Quantitative magnetic resonance immunohistochemistry with ligand-targeted F-19 nanoparticles, Magn. Reson. Med. 52(6):1255-1262 (2004)  CK3 MORAWSKI et al., Targeted Nanoparticles for Quantitative Imaging of Sparse Molecular Epitopes with MRK, Mag. Res. in Med. 51(3):480-486 (2004)  CL3 NEUBAUER et al., Endothelial stem cell detection in vivo with unique perfluorocarbon nanoparticle labels using fluorine (F-19) MNRI at 1.5 T, Circulation 114(18)(Suppl. S): 251 (Abstract) (2006)  CM3 PAKTLOW et al., Thelper 2 (Th2) T cells induce acute pancreatitis and diabetes in immune-compromised nonobese diabetic (NOD) mice, J. Exp. Med. 186(2):299-306 (1997)  CN3 PARTLOW et al., 19F magnetic resonance imaging for stem/progenitor cell tracking with multiple unique perfluorocarbon nanobeacons, FASEB J. 21:1647-1654 (2007)  CO3 PELCHEN-MATTHEWS et al., Phorbol ester-induced downregulation of CD4 is a multistep process involving dissociation from p56lok, increased association with clathrin-coated pits, and altered endosomal sorting, J. Exp. Med. 178(4):1209-1222 (1993)  CP3 PHILLIPS et al., MAdCAM-1 is needed for diabetes development mediated by the T cell clone, BDC-2.5, Immunology 116(4):525-531 (2005)  CQ3 PHILLIPS et al., Nondepleting anti-CD4 has an immediate action on diabetogenic effector cells, halting their destruction of pancreatic beta cells, J. Immunol. 165(4):1949-1955 (2000)  CR3 PIACENT1 et al., Synthesis and characterization of fluorinated polyetheric amides, J. Fluor. Chem. 68:227-235 (1994)  CS3 PINTASKE et al.	CF3	
diabetic non-obese diabetic (NOD) mice - a longitudinal-study, Clin. Exp. Immunol. 60(3):622-630 (1985)  CH3 MODO et al., Mapping transplanted stem cell migration after a stroke: a serial, in vivo magnetic resonance imaging study, Neuroimage 21(1):311-317 (2004)  CI3 MOORE et al., Tracking the recruitment of diabetogenic CD8+ T-cells to the pancreas in real time, Diabetes 53(6):1459-1466 (2004)  CJ3 MORAWSKI et al., Quantitative magnetic resonance immunohistochemistry with ligand-targeted F-19 nanoparticles, Magn. Reson. Med. 52(6):1255-1262 (2004)  CK3 MORAWSKI et al., Targeted Nanoparticles for Quantitative Imaging of Sparse Molecular Epitopes with MRK, Mag. Res. in Med. 51(3):480-486 (2004)  CL3 NEUBAUER et al., Endothelial stem cell detection in vivo with unique perflourocarbon nanoparticle labels using fluorine (F-19) MNRI at 1.5 T, Circulation 114(18)(Suppl. S): 251 (Abstract) (2006)  CM3 PAKALA et al., T helper 2 (Th2) T cells induce acute pancreatitis and diabetes in immune-compromised nonobese diabetic (NOD) mice, J. Exp. Med. 186(2):299-306 (1997)  CN3 PARTLOW et al., 19F magnetic resonance imaging for stem/progenitor cell tracking with multiple unique perfluorocarbon nanobeacons, FASEB J. 21:1647-1654 (2007)  CO3 PELCHEN-MATTHEWS et al., Phorbol ester-induced downregulation of CD4 is a multistep process involving dissociation from p56lck, increased association with clathrin-coated pits, and altered endosomal sorting, J. Exp. Med. 178(4):1209-1222 (1993)  CP3 PHILLIPS et al., MAdCAM-1 is needed for diabetes development mediated by the T cell clone, BDC-2.5, Immunology 116(4):525-531 (2005)  CQ3 PHILLIPS et al., Nondepleting anti-CD4 has an immediate action on diabetogenic effector cells, halting their destruction of pancreatic beta cells, J. Immunol. 165(4):1949-1955 (2000)  CR3 PIACENT1 et al., Synthesis and characterization of fluorinated polyetheric amides, J. Fluor. Chem. 68:227-235 (1994)  CS3 PINTASKE et al., In vivo dynamics of T cell activation, proliferation, and death in HIV-1 infection: why		
CH3 MODO et al., Mapping transplanted stem cell migration after a stroke: a serial, in vivo magnetic resonance imaging study, Neuroimage 21(1):311-317 (2004)  CI3 MOORE et al., Tracking the recruitment of diabetogenic CD8+ T-cells to the pancreas in real time, Diabetes 53(6):1459-1466 (2004)  CJ3 MORAWSKI et al., Quantitative magnetic resonance immunohistochemistry with ligand-targeted F-19 nanoparticles, Magn. Reson. Med. 52(6):1255-1262 (2004)  CK3 MORAWSKI et al., Targeted Nanoparticles for Quantitative Imaging of Sparse Molecular Epitopes with MRK, Mag. Res. in Med. 51(3):480-486 (2004)  CL3 NEUBAUER et al., Endothelial stem cell detection in vivo with unique perflourocarbon nanoparticle labels using fluorine (F-19) MNRI at 1.5 T, Circulation 114(18)(Suppl. S): 251 (Abstract) (2006)  CM3 PAKALA et al., T helper 2 (Th2) T cells induce acute pancreatitis and diabetes in immune-compromised nonobese diabetic (NOD) mice, J. Exp. Med. 186(2):299-306 (1997)  CN3 PARTLOW et al., 19F magnetic resonance imaging for stem/progenitor cell tracking with multiple unique perfluorocarbon nanobeacons, FASEB J. 21:1647-1654 (2007)  CO3 PELCHEN-MATTHEWS et al., Phorbol ester-induced downregulation of CD4 is a multistep process involving dissociation from p56lck, increased association with clathrin-coated pits, and altered endosomal sorting, J. Exp. Med. 178(4):1209-1222 (1993)  CP3 PHILLIPS et al., MAdCAM-1 is needed for diabetes development mediated by the T cell clone, BDC-2.5, Immunology 116(4):525-531 (2005)  CO3 PHILLIPS et al., Nondepleting anti-CD4 has an immediate action on diabetogenic effector cells, halting their destruction of pancreatic beta cells, J. Immunol. 165(4):1949-1955 (2000)  PARS PARS Res et al., A preparation technique for quantitative investigation of SPIO-containing solutions and SPIO-labelled cells by MRI, Biomed. Tech. 50(6):174-180 (2005) (English Abstract)  CT3 RIBEIRO et al., In vivo dynamics of T cell activation, proliferation, and death in HIV-1 infection: why are CD4+ but not CD8+ T cells depl	CG	
CH3 MODO et al., Mapping transplanted stem cell migration after a stroke: a serial, in vivo magnetic resonance imaging study, Neuroimage 21(1):311-317 (2004)  CI3 MOORE et al., Tracking the recruitment of diabetogenic CD8+ T-cells to the pancreas in real time, Diabetes 53(6):1459-1466 (2004)  CJ3 MORAWSKI et al., Quantitative magnetic resonance immunohistochemistry with ligand-targeted F-19 nanoparticles, Magn. Reson. Med. 52(6):1255-1262 (2004)  CK3 MORAWSKI et al., Targeted Nanoparticles for Quantitative Imaging of Sparse Molecular Epitopes with MRK, Mag. Res. in Med. 51(3):480-486 (2004)  CL3 NEUBAUER et al., Endothelial stem cell detection in vivo with unique perflourocarbon nanoparticle labels using fluorine (F-19) MNRI at 1.5 T, Circulation 114(18)(Suppl. S): 251 (Abstract) (2006)  CM3 PAKALA et al., T helper 2 (Th2) T cells induce acute pancreatitis and diabetes in immune-compromised nonobese diabetic (NOD) mice, J. Exp. Med. 186(2):299-306 (1997)  CN3 PARTLOW et al., 19F magnetic resonance imaging for stem/progenitor cell tracking with multiple unique perfluorocarbon nanobeacons, FASEB J. 21:1647-1654 (2007)  CO3 PELCHEN-MATTHEWS et al., Phorbol ester-induced downregulation of CD4 is a multistep process involving dissociation from p56lck, increased association with clathrin-coated pits, and altered endosomal sorting, J. Exp. Med. 178(4):1209-1222 (1993)  CP3 PHILLIPS et al., MAdCAM-1 is needed for diabetes development mediated by the T cell clone, BDC-2.5, Immunology 116(4):525-531 (2005)  CQ3 PHILLIPS et al., Nondepleting anti-CD4 has an immediate action on diabetogenic effector cells, halting their destruction of pancreatic beta cells, J. Immunol. 165(4):1949-1955 (2000)  PASE PASE Al., A preparation technique for quantitative investigation of SPIO-containing solutions and SPIO-labelled cells by MRI, Biomed. Tech. 50(6):174-180 (2005) (English Abstract)  RIBEIRO et al., In vivo dynamics of T cell activation, proliferation, and death in HIV-1 infection: why are CD4+ but not CD8+ T cells depleted? Proc.		
magnetic resonance imaging study, Neuroimage 21(1):311-317 (2004)  CI3 MOORE et al., Tracking the recruitment of diabetogenic CD8+ T-cells to the pancreas in real time, Diabetes 53(6):1459-1466 (2004)  CJ3 MORAWSKI et al., Quantitative magnetic resonance immunohistochemistry with ligand-targeted F-19 nanoparticles, Magn. Reson. Med. 52(6):1255-1262 (2004)  CK3 MORAWSKI et al., Targeted Nanoparticles for Quantitative Imaging of Sparse Molecular Epitopes with MRK, Mag. Res. in Med. 51(3):480-486 (2004)  CL3 NEUBAUER et al., Endothelial stem cell detection in vivo with unique perflourocarbon nanoparticle labels using fluorine (F-19) MNRI at 1.5 T, Circulation 114(18)(Suppl. S): 251 (Abstract) (2006)  CM3 PAKALA et al., Thelper 2 (Th2) T cells induce acute pancreatitis and diabetes in immune-compromised nonobese diabetic (NOD) mice, J. Exp. Med. 186(2):299-306 (1997)  CN3 PARTLOW et al., 19F magnetic resonance imaging for stem/progenitor cell tracking with multiple unique perfluorocarbon nanobeacons, FASEB J. 21:1647-1654 (2007)  CO3 PELCHEN-MATTHEWS et al., Phorbol ester-induced downregulation of CD4 is a multistep process involving dissociation from p56lck, increased association with clathrin-coated pits, and altered endosomal sorting, J. Exp. Med. 178(4):1209-1222 (1993)  CP3 PHILLIPS et al., MAdCAM-1 is needed for diabetes development mediated by the T cell clone, BDC-2.5, Immunology 116(4):525-531 (2005)  CQ3 PHILLIPS et al., Nondepleting anti-CD4 has an immediate action on diabetogenic effector cells, halting their destruction of pancreatic beta cells, J. Immunol. 165(4):1949-1955 (2000)  CR3 PIACENTI et al., Synthesis and characterization of fluorinated polyetheric amides, J. Fluor. Chem. 68:227-235 (1994)  CS3 PINTASKE et al., A preparation technique for quantitative investigation of SPIO-containing solutions and SPIO-labelled cells by MRI, Biomed. Tech. 50(6):174-180 (2005) (English Abstract)  CT3 RIBEIRO et al., In vivo dynamics of T cell activation, proliferation, and death in HIV-1 infection: why are CD		
CI3 MOORE et al., Tracking the recruitment of diabetogenic CD8+ T-cells to the pancreas in real time, Diabetes 53(6):1459-1466 (2004)  CJ3 MORAWSKI et al., Quantitative magnetic resonance immunohistochemistry with ligand-targeted F-19 nanoparticles, Magn. Reson. Med. 52(6):1255-1262 (2004)  CK3 MORAWSKI et al., Targeted Nanoparticles for Quantitative Imaging of Sparse Molecular Epitopes with MRK, Mag. Res. in Med. 51(3):480-486 (2004)  CL3 NEUBAUER et al., Endothelial stem cell detection in vivo with unique perflourocarbon nanoparticle labels using fluorine (F-19) MNRI at 1.5 T, Circulation 114(18)(Suppl. S): 251 (Abstract) (2006)  CM3 PAKALA et al., T helper 2 (Th2) T cells induce acute pancreatitis and diabetes in immune-compromised nonobese diabetic (NOD) mice, J. Exp. Med. 186(2):299-306 (1997)  CN3 PARTLOW et al., 19F magnetic resonance imaging for stem/progenitor cell tracking with multiple unique perfluorocarbon nanobeacons, FASEB J. 21:1647-1654 (2007)  CO3 PELCHEN-MATTHEWS et al., Phorbol ester-induced downregulation of CD4 is a multistep process involving dissociation from p56lck, increased association with clathrin-coated pits, and altered endosomal sorting, J. Exp. Med. 178(4):1209-1222 (1993)  CP3 PHILLIPS et al., MAdCAM-1 is needed for diabetes development mediated by the T cell clone, BDC-2.5, Immunology 116(4):525-531 (2005)  CQ3 PHILLIPS et al., Nondepleting anti-CD4 has an immediate action on diabetogenic effector cells, halting their destruction of pancreatic beta cells, J. Immunol. 165(4):1949-1955 (2000)  CR3 PIACENTI et al., Synthesis and characterization of fluorinated polyetheric amides, J. Fluor. Chem. 68:227-235 (1994)  CS3 PINTASKE et al., A preparation technique for quantitative investigation of SPIO-containing solutions and SPIO-labelled cells by MRI, Biomed. Tech. 50(6):174-180 (2005) (English Abstract)	CH3	
time, Diabetes 53(6):1459-1466 (2004)  CJ3 MORAWSKI et al., Quantitative magnetic resonance immunohistochemistry with ligand-targeted F-19 nanoparticles, Magn. Reson. Med. 52(6):1255-1262 (2004)  CK3 MORAWSKI et al., Targeted Nanoparticles for Quantitative Imaging of Sparse Molecular Epitopes with MRK, Mag. Res. in Med. 51(3):480-486 (2004)  CL3 NEUBAUER et al., Endothelial stem cell detection in vivo with unique perflourocarbon nanoparticle labels using fluorine (F-19) MNRI at 1.5 T, Circulation 114(18)(Suppl. S): 251 (Abstract) (2006)  CM3 PAKALA et al., T helper 2 (Th2) T cells induce acute pancreatitis and diabetes in immune-compromised nonobese diabetic (NOD) mice, J. Exp. Med. 186(2):299-306 (1997)  CN3 PARTLOW et al., 19F magnetic resonance imaging for stem/progenitor cell tracking with multiple unique perfluorocarbon nanobeacons, FASEB J. 21:1647-1654 (2007)  CO3 PELCHEN-MATTHEWS et al., Phorbol ester-induced downregulation of CD4 is a multistep process involving dissociation from p56lck, increased association with clathrin-coated pits, and altered endosomal sorting, J. Exp. Med. 178(4):1209-1222 (1993)  CP3 PHILLIPS et al., MAdCAM-1 is needed for diabetes development mediated by the T cell clone, BDC-2.5, Immunology 116(4):525-531 (2005)  CQ3 PHILLIPS et al., Nondepleting anti-CD4 has an immediate action on diabetogenic effector cells, halting their destruction of pancreatic beta cells, J. Immunol. 165(4):1949-1955 (2000)  CR3 PIACENTI et al., Synthesis and characterization of fluorinated polyetheric amides, J. Fluor. Chem. 68:227-235 (1994)  CS3 PINTASKE et al., A preparation technique for quantitative investigation of SPIO-containing solutions and SPIO-labelled cells by MRI, Biomed. Tech. 50(6):174-180 (2005) (English Abstract)		
CJ3 MORAWSKI et al., Quantitative magnetic resonance immunohistochemistry with ligand-targeted F-19 nanoparticles, Magn. Reson. Med. 52(6):1255-1262 (2004)  CK3 MORAWSKI et al., Targeted Nanoparticles for Quantitative Imaging of Sparse Molecular Epitopes with MRK, Mag. Res. in Med. 51(3):480-486 (2004)  CL3 NEUBAUER et al., Endothelial stem cell detection in vivo with unique perflourocarbon nanoparticle labels using fluorine (F-19) MNRI at 1.5 T, Circulation 114(18)(Suppl. S): 251 (Abstract) (2006)  CM3 PAKALA et al., T helper 2 (Th2) T cells induce acute pancreatitis and diabetes in immune-compromised nonobese diabetic (NOD) mice, J. Exp. Med. 186(2):299-306 (1997)  CN3 PARTLOW et al., 19F magnetic resonance imaging for stem/progenitor cell tracking with multiple unique perfluorocarbon nanobeacons, FASEB J. 21:1647-1654 (2007)  CO3 PELCHEN-MATTHEWS et al., Phorbol ester-induced downregulation of CD4 is a multistep process involving dissociation from p56lck, increased association with clathrin-coated pits, and altered endosomal sorting, J. Exp. Med. 178(4):1209-1222 (1993)  CP3 PHILLIPS et al., MAdCAM-1 is needed for diabetes development mediated by the T cell clone, BDC-2.5, Immunology 116(4):525-531 (2005)  PHILLIPS et al., Nondepleting anti-CD4 has an immediate action on diabetogenic effector cells, halting their destruction of pancreatic beta cells, J. Immunol. 165(4):1949-1955 (2000)  PIACENTI et al., Synthesis and characterization of fluorinated polyetheric amides, J. Fluor. Chem. 68:227-235 (1994)  CS3 PINTASKE et al., A preparation technique for quantitative investigation of SPIO-containing solutions and SPIO-labelled cells by MRI, Biomed. Tech. 50(6):174-180 (2005) (English Abstract)  CT3 RIBEIRO et al., In vivo dynamics of T cell activation, proliferation, and death in HIV-1 infection: why are CD4+ but not CD8+ T cells depleted? Proc. Natl. Acad. Sci. USA 99(24):15572-15577	Cl3	
CK3 MORAWSKI et al., Targeted Nanoparticles for Quantitative Imaging of Sparse Molecular Epitopes with MRK, Mag. Res. in Med. 51(3):480-486 (2004)  CL3 NEUBAUER et al., Endothelial stem cell detection in vivo with unique perflourocarbon nanoparticle labels using fluorine (F-19) MNRI at 1.5 T, Circulation 114(18)(Suppl. S): 251 (Abstract) (2006)  CM3 PAKALA et al., T helper 2 (Th2) T cells induce acute pancreatitis and diabetes in immune-compromised nonobese diabetic (NOD) mice, J. Exp. Med. 186(2):299-306 (1997)  CN3 PARTLOW et al., 19F magnetic resonance imaging for stem/progenitor cell tracking with multiple unique perfluorocarbon nanobeacons, FASEB J. 21:1647-1654 (2007)  CO3 PELCHEN-MATTHEWS et al., Phorbol ester-induced downregulation of CD4 is a multistep process involving dissociation from p56lck, increased association with clathrin-coated pits, and altered endosomal sorting, J. Exp. Med. 178(4):1209-1222 (1993)  CP3 PHILLIPS et al., MAdCAM-1 is needed for diabetes development mediated by the T cell clone, BDC-2.5, Immunology 116(4):525-531 (2005)  CQ3 PHILLIPS et al., Nondepleting anti-CD4 has an immediate action on diabetogenic effector cells, halting their destruction of pancreatic beta cells, J. Immunol. 165(4):1949-1955 (2000)  CR3 PIACENTI et al., Synthesis and characterization of fluorinated polyetheric amides, J. Fluor. Chem. 68:227-235 (1994)  CS3 PINTASKE et al., A preparation technique for quantitative investigation of SPIO-containing solutions and SPIO-labelled cells by MRI, Biomed. Tech. 50(6):174-180 (2005) (English Abstract)  CT3 RIBEIRO et al., In vivo dynamics of T cell activation, proliferation, and death in HIV-1 infection: why are CD4+ but not CD8+ T cells depleted? Proc. Natl. Acad. Sci. USA 99(24):15572-15577	CJ3	MORAWSKI et al., Quantitative magnetic resonance immunohistochemistry with ligand-
CK3 MORAWSKI et al., Targeted Nanoparticles for Quantitative Imaging of Sparse Molecular Epitopes with MRK, Mag. Res. in Med. 51(3):480-486 (2004)  CL3 NEUBAUER et al., Endothelial stem cell detection in vivo with unique perflourocarbon nanoparticle labels using fluorine (F-19) MNRI at 1.5 T, Circulation 114(18)(Suppl. S): 251 (Abstract) (2006)  CM3 PAKALA et al., T helper 2 (Th2) T cells induce acute pancreatitis and diabetes in immune-compromised nonobese diabetic (NOD) mice, J. Exp. Med. 186(2):299-306 (1997)  CN3 PARTLOW et al., 19F magnetic resonance imaging for stem/progenitor cell tracking with multiple unique perfluorocarbon nanobeacons, FASEB J. 21:1647-1654 (2007)  CO3 PELCHEN-MATTHEWS et al., Phorbol ester-induced downregulation of CD4 is a multistep process involving dissociation from p56lck, increased association with clathrin-coated pits, and altered endosomal sorting, J. Exp. Med. 178(4):1209-1222 (1993)  CP3 PHILLIPS et al., MAdCAM-1 is needed for diabetes development mediated by the T cell clone, BDC-2.5, Immunology 116(4):525-531 (2005)  CQ3 PHILLIPS et al., Nondepleting anti-CD4 has an immediate action on diabetogenic effector cells, halting their destruction of pancreatic beta cells, J. Immunol. 165(4):1949-1955 (2000)  CR3 PIACENTI et al., Synthesis and characterization of fluorinated polyetheric amides, J. Fluor. Chem. 68:227-235 (1994)  CS3 PINTASKE et al., A preparation technique for quantitative investigation of SPIO-containing solutions and SPIO-labelled cells by MRI, Biomed. Tech. 50(6):174-180 (2005) (English Abstract)  CT3 RIBEIRO et al., In vivo dynamics of T cell activation, proliferation, and death in HIV-1 infection: why are CD4+ but not CD8+ T cells depleted? Proc. Natl. Acad. Sci. USA 99(24):15572-15577		targeted F-19 nanoparticles, Magn. Reson. Med. 52(6):1255-1262 (2004)
CL3 NEUBAUER et al., Endothelial stem cell detection in vivo with unique perflourocarbon nanoparticle labels using fluorine (F-19) MNRI at 1.5 T, Circulation 114(18)(Suppl. S): 251 (Abstract) (2006)  CM3 PAKALA et al., T helper 2 (Th2) T cells induce acute pancreatitis and diabetes in immune-compromised nonobese diabetic (NOD) mice, J. Exp. Med. 186(2):299-306 (1997)  CN3 PARTLOW et al., 19F magnetic resonance imaging for stem/progenitor cell tracking with multiple unique perfluorocarbon nanobeacons, FASEB J. 21:1647-1654 (2007)  CO3 PELCHEN-MATTHEWS et al., Phorbol ester-induced downregulation of CD4 is a multistep process involving dissociation from p56lck, increased association with clathrin-coated pits, and altered endosomal sorting, J. Exp. Med. 178(4):1209-1222 (1993)  CP3 PHILLIPS et al., MAdCAM-1 is needed for diabetes development mediated by the T cell clone, BDC-2.5, Immunology 116(4):525-531 (2005)  CQ3 PHILLIPS et al., Nondepleting anti-CD4 has an immediate action on diabetogenic effector cells, halting their destruction of pancreatic beta cells, J. Immunol. 165(4):1949-1955 (2000)  CR3 PIACENTI et al., Synthesis and characterization of fluorinated polyetheric amides, J. Fluor. Chem. 68:227-235 (1994)  CS3 PINTASKE et al., A preparation technique for quantitative investigation of SPIO-containing solutions and SPIO-labelled cells by MRI, Biomed. Tech. 50(6):174-180 (2005) (English Abstract)  CT3 RIBEIRO et al., In vivo dynamics of T cell activation, proliferation, and death in HIV-1 infection: why are CD4+ but not CD8+ T cells depleted? Proc. Natl. Acad. Sci. USA 99(24):15572-15577	CK3	MORAWSKI et al., Targeted Nanoparticles for Quantitative Imaging of Sparse Molecular
nanoparticle labels using fluorine (F-19) MNRI at 1.5 T, Circulation 114(18)(Suppl. S): 251 (Abstract) (2006)  CM3 PAKALA et al., T helper 2 (Th2) T cells induce acute pancreatitis and diabetes in immune-compromised nonobese diabetic (NOD) mice, J. Exp. Med. 186(2):299-306 (1997)  CN3 PARTLOW et al., 19F magnetic resonance imaging for stem/progenitor cell tracking with multiple unique perfluorocarbon nanobeacons, FASEB J. 21:1647-1654 (2007)  CO3 PELCHEN-MATTHEWS et al., Phorbol ester-induced downregulation of CD4 is a multistep process involving dissociation from p56lck, increased association with clathrin-coated pits, and altered endosomal sorting, J. Exp. Med. 178(4):1209-1222 (1993)  CP3 PHILLIPS et al., MAdCAM-1 is needed for diabetes development mediated by the T cell clone, BDC-2.5, Immunology 116(4):525-531 (2005)  CQ3 PHILLIPS et al., Nondepleting anti-CD4 has an immediate action on diabetogenic effector cells, halting their destruction of pancreatic beta cells, J. Immunol. 165(4):1949-1955 (2000)  CR3 PIACENTI et al., Synthesis and characterization of fluorinated polyetheric amides, J. Fluor. Chem. 68:227-235 (1994)  CS3 PINTASKE et al., A preparation technique for quantitative investigation of SPIO-containing solutions and SPIO-labelled cells by MRI, Biomed. Tech. 50(6):174-180 (2005) (English Abstract)  CT3 RIBEIRO et al., In vivo dynamics of T cell activation, proliferation, and death in HIV-1 infection: why are CD4+ but not CD8+ T cells depleted? Proc. Natl. Acad. Sci. USA 99(24):15572-15577		
(Abstract) (2006)  CM3 PAKALA et al., T helper 2 (Th2) T cells induce acute pancreatitis and diabetes in immune-compromised nonobese diabetic (NOD) mice, J. Exp. Med. 186(2):299-306 (1997)  CN3 PARTLOW et al., 19F magnetic resonance imaging for stem/progenitor cell tracking with multiple unique perfluorocarbon nanobeacons, FASEB J. 21:1647-1654 (2007)  CO3 PELCHEN-MATTHEWS et al., Phorbol ester-induced downregulation of CD4 is a multistep process involving dissociation from p56lck, increased association with clathrin-coated pits, and altered endosomal sorting, J. Exp. Med. 178(4):1209-1222 (1993)  CP3 PHILLIPS et al., MAdCAM-1 is needed for diabetes development mediated by the T cell clone, BDC-2.5, Immunology 116(4):525-531 (2005)  CQ3 PHILLIPS et al., Nondepleting anti-CD4 has an immediate action on diabetogenic effector cells, halting their destruction of pancreatic beta cells, J. Immunol. 165(4):1949-1955 (2000)  CR3 PIACENTI et al., Synthesis and characterization of fluorinated polyetheric amides, J. Fluor. Chem. 68:227-235 (1994)  CS3 PINTASKE et al., A preparation technique for quantitative investigation of SPIO-containing solutions and SPIO-labelled cells by MRI, Biomed. Tech. 50(6):174-180 (2005) (English Abstract)  CT3 RIBEIRO et al., In vivo dynamics of T cell activation, proliferation, and death in HIV-1 infection: why are CD4+ but not CD8+ T cells depleted? Proc. Natl. Acad. Sci. USA 99(24):15572-15577	CL3	
CM3 PAKALA et al., T helper 2 (Th2) T cells induce acute pancreatitis and diabetes in immune-compromised nonobese diabetic (NOD) mice, J. Exp. Med. 186(2):299-306 (1997)  CN3 PARTLOW et al., 19F magnetic resonance imaging for stem/progenitor cell tracking with multiple unique perfluorocarbon nanobeacons, FASEB J. 21:1647-1654 (2007)  CO3 PELCHEN-MATTHEWS et al., Phorbol ester-induced downregulation of CD4 is a multistep process involving dissociation from p56lck, increased association with clathrin-coated pits, and altered endosomal sorting, J. Exp. Med. 178(4):1209-1222 (1993)  CP3 PHILLIPS et al., MAdCAM-1 is needed for diabetes development mediated by the T cell clone, BDC-2.5, Immunology 116(4):525-531 (2005)  CQ3 PHILLIPS et al., Nondepleting anti-CD4 has an immediate action on diabetogenic effector cells, halting their destruction of pancreatic beta cells, J. Immunol. 165(4):1949-1955 (2000)  CR3 PIACENTI et al., Synthesis and characterization of fluorinated polyetheric amides, J. Fluor. Chem. 68:227-235 (1994)  CS3 PINTASKE et al., A preparation technique for quantitative investigation of SPIO-containing solutions and SPIO-labelled cells by MRI, Biomed. Tech. 50(6):174-180 (2005) (English Abstract)  CT3 RIBEIRO et al., In vivo dynamics of T cell activation, proliferation, and death in HIV-1 infection: why are CD4+ but not CD8+ T cells depleted? Proc. Natl. Acad. Sci. USA 99(24):15572-15577		
compromised nonobese diabetic (NOD) mice, J. Exp. Med. 186(2):299-306 (1997)  CN3 PARTLOW et al., 19F magnetic resonance imaging for stem/progenitor cell tracking with multiple unique perfluorocarbon nanobeacons, FASEB J. 21:1647-1654 (2007)  CO3 PELCHEN-MATTHEWS et al., Phorbol ester-induced downregulation of CD4 is a multistep process involving dissociation from p56lck, increased association with clathrin-coated pits, and altered endosomal sorting, J. Exp. Med. 178(4):1209-1222 (1993)  CP3 PHILLIPS et al., MAdCAM-1 is needed for diabetes development mediated by the T cell clone, BDC-2.5, Immunology 116(4):525-531 (2005)  CQ3 PHILLIPS et al., Nondepleting anti-CD4 has an immediate action on diabetogenic effector cells, halting their destruction of pancreatic beta cells, J. Immunol. 165(4):1949-1955 (2000)  CR3 PIACENTI et al., Synthesis and characterization of fluorinated polyetheric amides, J. Fluor. Chem. 68:227-235 (1994)  CS3 PINTASKE et al., A preparation technique for quantitative investigation of SPIO-containing solutions and SPIO-labelled cells by MRI, Biomed. Tech. 50(6):174-180 (2005) (English Abstract)  CT3 RIBEIRO et al., In vivo dynamics of T cell activation, proliferation, and death in HIV-1 infection: why are CD4+ but not CD8+ T cells depleted? Proc. Natl. Acad. Sci. USA 99(24):15572-15577		
CN3 PARTLOW et al., 19F magnetic resonance imaging for stem/progenitor cell tracking with multiple unique perfluorocarbon nanobeacons, FASEB J. 21:1647-1654 (2007)  CO3 PELCHEN-MATTHEWS et al., Phorbol ester-induced downregulation of CD4 is a multistep process involving dissociation from p56lck, increased association with clathrin-coated pits, and altered endosomal sorting, J. Exp. Med. 178(4):1209-1222 (1993)  CP3 PHILLIPS et al., MAdCAM-1 is needed for diabetes development mediated by the T cell clone, BDC-2.5, Immunology 116(4):525-531 (2005)  CQ3 PHILLIPS et al., Nondepleting anti-CD4 has an immediate action on diabetogenic effector cells, halting their destruction of pancreatic beta cells, J. Immunol. 165(4):1949-1955 (2000)  CR3 PIACENTI et al., Synthesis and characterization of fluorinated polyetheric amides, J. Fluor. Chem. 68:227-235 (1994)  CS3 PINTASKE et al., A preparation technique for quantitative investigation of SPIO-containing solutions and SPIO-labelled cells by MRI, Biomed. Tech. 50(6):174-180 (2005) (English Abstract)  RIBEIRO et al., In vivo dynamics of T cell activation, proliferation, and death in HIV-1 infection: why are CD4+ but not CD8+ T cells depleted? Proc. Natl. Acad. Sci. USA 99(24):15572-15577	CM	
multiple unique perfluorocarbon nanobeacons, FASEB J. 21:1647-1654 (2007)  CO3 PELCHEN-MATTHEWS et al., Phorbol ester-induced downregulation of CD4 is a multistep process involving dissociation from p56lck, increased association with clathrin-coated pits, and altered endosomal sorting, J. Exp. Med. 178(4):1209-1222 (1993)  CP3 PHILLIPS et al., MAdCAM-1 is needed for diabetes development mediated by the T cell clone, BDC-2.5, Immunology 116(4):525-531 (2005)  CQ3 PHILLIPS et al., Nondepleting anti-CD4 has an immediate action on diabetogenic effector cells, halting their destruction of pancreatic beta cells, J. Immunol. 165(4):1949-1955 (2000)  CR3 PIACENTI et al., Synthesis and characterization of fluorinated polyetheric amides, J. Fluor. Chem. 68:227-235 (1994)  CS3 PINTASKE et al., A preparation technique for quantitative investigation of SPIO-containing solutions and SPIO-labelled cells by MRI, Biomed. Tech. 50(6):174-180 (2005) (English Abstract)  CT3 RIBEIRO et al., In vivo dynamics of T cell activation, proliferation, and death in HIV-1 infection: why are CD4+ but not CD8+ T cells depleted? Proc. Natl. Acad. Sci. USA 99(24):15572-15577		
CO3 PELCHEN-MATTHEWS et al., Phorbol ester-induced downregulation of CD4 is a multistep process involving dissociation from p56lck, increased association with clathrin-coated pits, and altered endosomal sorting, J. Exp. Med. 178(4):1209-1222 (1993)  CP3 PHILLIPS et al., MAdCAM-1 is needed for diabetes development mediated by the T cell clone, BDC-2.5, Immunology 116(4):525-531 (2005)  CQ3 PHILLIPS et al., Nondepleting anti-CD4 has an immediate action on diabetogenic effector cells, halting their destruction of pancreatic beta cells, J. Immunol. 165(4):1949-1955 (2000)  CR3 PIACENTI et al., Synthesis and characterization of fluorinated polyetheric amides, J. Fluor. Chem. 68:227-235 (1994)  CS3 PINTASKE et al., A preparation technique for quantitative investigation of SPIO-containing solutions and SPIO-labelled cells by MRI, Biomed. Tech. 50(6):174-180 (2005) (English Abstract)  CT3 RIBEIRO et al., In vivo dynamics of T cell activation, proliferation, and death in HIV-1 infection: why are CD4+ but not CD8+ T cells depleted? Proc. Natl. Acad. Sci. USA 99(24):15572-15577	CN3	
process involving dissociation from p56lck, increased association with clathrin-coated pits, and altered endosomal sorting, J. Exp. Med. 178(4):1209-1222 (1993)  CP3 PHILLIPS et al., MAdCAM-1 is needed for diabetes development mediated by the T cell clone, BDC-2.5, Immunology 116(4):525-531 (2005)  CQ3 PHILLIPS et al., Nondepleting anti-CD4 has an immediate action on diabetogenic effector cells, halting their destruction of pancreatic beta cells, J. Immunol. 165(4):1949-1955 (2000)  CR3 PIACENTI et al., Synthesis and characterization of fluorinated polyetheric amides, J. Fluor. Chem. 68:227-235 (1994)  CS3 PINTASKE et al., A preparation technique for quantitative investigation of SPIO-containing solutions and SPIO-labelled cells by MRI, Biomed. Tech. 50(6):174-180 (2005) (English Abstract)  CT3 RIBEIRO et al., In vivo dynamics of T cell activation, proliferation, and death in HIV-1 infection: why are CD4+ but not CD8+ T cells depleted? Proc. Natl. Acad. Sci. USA 99(24):15572-15577		
altered endosomal sorting, J. Exp. Med. 178(4):1209-1222 (1993)  CP3 PHILLIPS et al., MAdCAM-1 is needed for diabetes development mediated by the T cell clone, BDC-2.5, Immunology 116(4):525-531 (2005)  CQ3 PHILLIPS et al., Nondepleting anti-CD4 has an immediate action on diabetogenic effector cells, halting their destruction of pancreatic beta cells, J. Immunol. 165(4):1949-1955 (2000)  CR3 PIACENTI et al., Synthesis and characterization of fluorinated polyetheric amides, J. Fluor. Chem. 68:227-235 (1994)  CS3 PINTASKE et al., A preparation technique for quantitative investigation of SPIO-containing solutions and SPIO-labelled cells by MRI, Biomed. Tech. 50(6):174-180 (2005) (English Abstract)  CT3 RIBEIRO et al., In vivo dynamics of T cell activation, proliferation, and death in HIV-1 infection: why are CD4+ but not CD8+ T cells depleted? Proc. Natl. Acad. Sci. USA 99(24):15572-15577		
CP3 PHILLIPS et al., MAdCAM-1 is needed for diabetes development mediated by the T cell clone, BDC-2.5, Immunology 116(4):525-531 (2005)  CQ3 PHILLIPS et al., Nondepleting anti-CD4 has an immediate action on diabetogenic effector cells, halting their destruction of pancreatic beta cells, J. Immunol. 165(4):1949-1955 (2000)  CR3 PIACENTI et al., Synthesis and characterization of fluorinated polyetheric amides, J. Fluor. Chem. 68:227-235 (1994)  CS3 PINTASKE et al., A preparation technique for quantitative investigation of SPIO-containing solutions and SPIO-labelled cells by MRI, Biomed. Tech. 50(6):174-180 (2005) (English Abstract)  CT3 RIBEIRO et al., In vivo dynamics of T cell activation, proliferation, and death in HIV-1 infection: why are CD4+ but not CD8+ T cells depleted? Proc. Natl. Acad. Sci. USA 99(24):15572-15577		
BDC-2.5, Immunology 116(4):525-531 (2005)  CQ3 PHILLIPS et al., Nondepleting anti-CD4 has an immediate action on diabetogenic effector cells, halting their destruction of pancreatic beta cells, J. Immunol. 165(4):1949-1955 (2000)  CR3 PIACENTI et al., Synthesis and characterization of fluorinated polyetheric amides, J. Fluor. Chem. 68:227-235 (1994)  CS3 PINTASKE et al., A preparation technique for quantitative investigation of SPIO-containing solutions and SPIO-labelled cells by MRI, Biomed. Tech. 50(6):174-180 (2005) (English Abstract)  CT3 RIBEIRO et al., In vivo dynamics of T cell activation, proliferation, and death in HIV-1 infection: why are CD4+ but not CD8+ T cells depleted? Proc. Natl. Acad. Sci. USA 99(24):15572-15577		
CQ3 PHILLIPS et al., Nondepleting anti-CD4 has an immediate action on diabetogenic effector cells, halting their destruction of pancreatic beta cells, J. Immunol. 165(4):1949-1955 (2000)  CR3 PIACENTI et al., Synthesis and characterization of fluorinated polyetheric amides, J. Fluor. Chem. 68:227-235 (1994)  CS3 PINTASKE et al., A preparation technique for quantitative investigation of SPIO-containing solutions and SPIO-labelled cells by MRI, Biomed. Tech. 50(6):174-180 (2005) (English Abstract)  CT3 RIBEIRO et al., In vivo dynamics of T cell activation, proliferation, and death in HIV-1 infection: why are CD4+ but not CD8+ T cells depleted? Proc. Natl. Acad. Sci. USA 99(24):15572-15577		
cells, halting their destruction of pancreatic beta cells, J. Immunol. 165(4):1949-1955 (2000)  CR3 PIACENTI et al., Synthesis and characterization of fluorinated polyetheric amides, J. Fluor. Chem. 68:227-235 (1994)  CS3 PINTASKE et al., A preparation technique for quantitative investigation of SPIO-containing solutions and SPIO-labelled cells by MRI, Biomed. Tech. 50(6):174-180 (2005) (English Abstract)  CT3 RIBEIRO et al., In vivo dynamics of T cell activation, proliferation, and death in HIV-1 infection: why are CD4+ but not CD8+ T cells depleted? Proc. Natl. Acad. Sci. USA 99(24):15572-15577	CO3	
CR3 PIACENTI et al., Synthesis and characterization of fluorinated polyetheric amides, J. Fluor. Chem. 68:227-235 (1994)  CS3 PINTASKE et al., A preparation technique for quantitative investigation of SPIO-containing solutions and SPIO-labelled cells by MRI, Biomed. Tech. 50(6):174-180 (2005) (English Abstract)  CT3 RIBEIRO et al., In vivo dynamics of T cell activation, proliferation, and death in HIV-1 infection: why are CD4+ but not CD8+ T cells depleted? Proc. Natl. Acad. Sci. USA 99(24):15572-15577		
Chem. 68:227-235 (1994)  CS3 PINTASKE et al., A preparation technique for quantitative investigation of SPIO-containing solutions and SPIO-labelled cells by MRI, Biomed. Tech. 50(6):174-180 (2005) (English Abstract)  CT3 RIBEIRO et al., In vivo dynamics of T cell activation, proliferation, and death in HIV-1 infection: why are CD4+ but not CD8+ T cells depleted? Proc. Natl. Acad. Sci. USA 99(24):15572-15577	CRS	
solutions and SPIO-labelled cells by MRI, Biomed. Tech. 50(6):174-180 (2005) (English Abstract)  CT3 RIBEIRO et al., In vivo dynamics of T cell activation, proliferation, and death in HIV-1 infection: why are CD4+ but not CD8+ T cells depleted? Proc. Natl. Acad. Sci. USA 99(24):15572-15577		Chem. 68:227-235 (1994)
Abstract)  CT3 RIBEIRO et al., In vivo dynamics of T cell activation, proliferation, and death in HIV-1 infection: why are CD4+ but not CD8+ T cells depleted? Proc. Natl. Acad. Sci. USA 99(24):15572-15577	CS	PINTASKE et al., A preparation technique for quantitative investigation of SPIO-containing
CT3 RIBEIRO et al., In vivo dynamics of T cell activation, proliferation, and death in HIV-1 infection: why are CD4+ but not CD8+ T cells depleted? Proc. Natl. Acad. Sci. USA 99(24):15572-15577		solutions and SPIO-labelled cells by MRI, Biomed. Tech. 50(6):174-180 (2005) (English
why are CD4+ but not CD8+ T cells depleted? Proc. Natl. Acad. Sci. USA 99(24):15572-15577		
	CT3	
(2002)		
		(2002)

Complete if Known Substitute for form 1449/PTO 10/586,015 Application Number **INFORMATION DISCLOSURE** August 4, 2008 Filing Date STATEMENT BY APPLICANT First Named Inventor Eric T. Ahrens Art Unit 1618 (Use as many sheets as necessary) Jagadishwar Rao Samala Examiner Name 4 of 4 CAMU-P01-002 Sheet Attorney Docket Number

0110	DODDIOUET 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1	
CO3	RODRIGUEZ et al., In vitro characterization of an Fe(8) cluster as potential MRI contrast	
	agent, NMR Biomed. 18(5):300-307 (2005)	
CV3	SCHNEIDER et al., In vivo microscopic evaluation of the microvascular behavior of FITC-	
	labeled macromolecular MR contrast agents in the hamster skinfold chamber, Invest. Radiol.	
	35(9):564-570 (2000)	
CW3	SHAPIRO et al., In vivo detection of single cells by MRI, Magn. Reson. Med. 55(2):242-249	
	(2006)	
CX3	SOLOSKI, Synthesis of perfluoro (polyether) difunctional compounds, J. Fluor. Chem. 11:601-	
	612 (1978)	
CY3	SOTAK et al., A new perfluorocarbon for use on fluorine-19 magnetic resonance imaging and	
	spectroscopy, Magn. Reson. Med. 29:188 (1993)	
CZ3	SRINIVAS et al., Fluorine-19 MRI for visualization and quantification of cell migration in a	
020	diabetes model, Mag. Res. In Med. 58(4):725-734 (2007)	
CA4	TAYLOR and DEUTSCH, 19F-nuclear magnetic resonance: measurements of [O2] and pH in	
	biological systems, Biophys J. 53: 227-233 (1988)	
CB4	TONELLI et al., Linear perfluoropolyether difunctional oligomers: chemistry, properties and	
CD4		
004	applications, J. Fluorine Chem. 95:51-70 (1999)	
CC4	TONELLI et al., Perfluoropolyether alkyl diesters: Structure effects of the alkyl group on the	
	kinetics of the hydrolysis reactions, J. Polym. Sci. Part A: Polym Chem. 40:4266-4280 (2002)	
CD4	TONELLI et al., Perfluoropolyether functional oligomers: unusual reactivity in organic	
	chemistry, J. Fluor. Chem. 118(1-2):107-121 (2002)	
CE4	TURVEY et al., Noninvasive imaging of pancreatic inflammation and its reversal in type 1	
	diabetes, J. Clin. Invest. 115(9):2454-2461 (2005)	
CF4	VENANZI et al., Structural properties and photophysical behavior of conformationally	
	constrained hexapeptides functionalized with a new fluorescent analog of tryptophan and a	
	nitroxide radical quencher, Biopolymers 75(2):128-139 (2004)	
CG4	WILSON et al., Measurement of preretinal oxygen-tension in the vitrectomized human eye	
	using F-19 magnetic resonance spectroscopy, Arch. Ophthalmol-Chic. 110(8):1098-1100	
	(1992)	
CH4	WISNER et al., A modular lymphographic magnetic resonance imaging contrast agent:	
	contrast enhancement with DNA transfection potential, J. Med. Chem. 40(25):3992-3996	
	(1997)	
CI4	WU et al., In situ labeling of immune cells with iron oxide particles: An approach to detect	
	organ rejection by cellular MRI, Proc. Natl. Acad. Sci. USA 103(6):1852-1857 (2006)	
CJ4	XIA et al., Tumour oxygen dynamics measured simultaneously by near-infrared spectroscopy	
557	and F-19 magnetic resonance imaging in rats, Phys. Med. Biol. 51(1):45-60 (2006)	
CK4	YEH et al., Intracellular labeling of T-cells with superparamagnetic contrast agents, Magn.	
	Reson. Med. 30(5):617-625 (1993)	
CL4	YOU et al., Detection and characterization of T cells specific for BDC2.5 T cell-stimulating	
CNAA	peptides, J. Immunol. 170(8):4011-4020 (2003)  YU et al., High-resolution MRI characterization of human thrombus using a novel fibrin-	
CM4		
0004	targeted paramagnetic nanoparticle contrast agent, Mag. Res. In Med. 44:867-872 (2000)	
CN4	ZHANG et al., Synthetic applications of fluorous solid-phase extraction (F-SPE), Tetrahedron	
	62:11837-11865 (2006)	

Examiner	Date	
Signature	Considered	

<sup>\*</sup>EXAMINER: Initial if reference considered, whether or not citation is in conformance with MPEP 609. Draw line through citation if not in conformance and not considered. Include copy of this form with next communication to applicant.

<sup>&</sup>lt;sup>1</sup>Applicant's unique citation designation number (optional). <sup>2</sup>Applicant is to place a check mark here if English language Translation is attached.